arXiv:math/0308113v1 [math.AC] 12 Aug 2003

GENERIC COHEN-MACAULAY MONOMIAL IDEALS

ABDUL SALAM JARRAH AND REINHARD LAUBENBACHER

Abstract. Given a simplicial complex, it is easy to construct a generic deformation of its Stanley-Reisner ideal. The main question under investigation in this paper is how to characterize the simplicial complexes such that their Stanley-Reisner ideals have Cohen-Macaulay generic deformations. Algorithms are presented to construct such deformations for matroid complexes, shifted complexes, and tree complexes.

1. Introduction

Let $S := k[x_1, \dots, x_n]$ be a polynomial ring in n variables over a field k. A monomial $\mathbf{x}^{\mathbf{a}} := x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n}$ is called *square-free* if $a_i = 0$ or 1 for all $1 \leq i \leq n$. For monomials $\mathbf{x}^{\mathbf{a}}$ and $\mathbf{x}^{\mathbf{b}}$, we say $\mathbf{x}^{\mathbf{a}}$ *strictly divides* $\mathbf{x}^{\mathbf{b}}$ if, for $1 \leq i \leq n$ and $a_i > 0$, we have $a_i < b_i$. A *monomial ideal* M is an ideal minimally generated by monomials, say $m_1, \dots, m_t$, (written as $M = \langle m_1, \dots, m_t \rangle$). The monomial ideal M is called *square-free* if all m_i are square free monomials.

Definition 1.1. A monomial ideal $M = \langle m_1, \dots, m_t \rangle$ is called *generic* if for any two distinct minimal generators m_i and m_j with the same positive degree in some variable x_h, there exists a third minimal generator m_l such that m_l strictly divides $\operatorname{lcm}(m_i, m_j)$.

Example 1.2. The monomial ideal $\langle x^2y^2, x^2z^2, xz \rangle \subset k[x, y, z]$ is generic but $\langle xy^2, xz^2, xz \rangle \subset k[x, y, z]$ is not generic.

This definition of generic monomial ideals has appeared in [5], generalizing a definition in [1]. See [10] for more information about generic monomial ideals.

Definition 1.3. A simplicial complex Γ on the vertex set $[n] := \{1, \dots, n\}$ is a non-empty collection of subsets of $[n]$ such that $\{i\} \in \Gamma$ for all $i \in [n]$, and whenever $\tau \subset \sigma$ and $\sigma \in \Gamma$, then $\tau \in \Gamma$. The elements of Γ are called *faces*, and the *dimension* of the face $\sigma \in \Gamma$ is $\dim(\sigma) := |\sigma| - 1$. The dimension of Γ is $\dim(\Gamma) := \max\{\dim(\sigma) : \sigma \in \Gamma\}$. Faces of dimension zero are called *vertices* and faces that are maximal under inclusion are called *facets*. The

---

Date: February 1, 2008.

1991 *Mathematics Subject Classification.* Primary , 13P10, 13C14, 52B20 ; Secondary 52B40, 52B22.

Key words and phrases. Monomial ideals, simplicial complexes, Cohen-Macaulay.

set of *minimal non-faces* of Γ is $\Sigma_\Gamma := \{\sigma \subseteq [n] : \sigma \notin \Gamma \text{ but } \sigma \setminus \{i\} \in \Gamma \text{ for each } i \in \sigma\}$.

There is a one-to-one correspondence between square-free monomial ideals in S and simplicial complexes on $[n]$. Given a simplicial complex Γ on the vertex set $[n]$, the *Stanley-Reisner* ideal I_Γ of Γ is the square-free monomial ideal $\langle x_{i_1} \cdots x_{i_s} : \{i_1, \ldots, i_s\} \in [n] \setminus \Gamma\rangle \subset S$. Conversely, for a given non-zero square-free monomial ideal $M \subset S$, there exists a simplicial complex Γ on $[n]$ such that $I_\Gamma = M$.

Definition 1.4. Let $I = \langle m_1, \ldots, m_t\rangle$ be a square-free monomial ideal. A *deformation* of I is a monomial ideal $M = \langle m_1^*, \ldots, m_t^*\rangle$ such that, for all $1 \leq i \leq t$, $m_i^* = x_{i_1}^{j_1} \cdots x_{i_s}^{j_s}$, where $m_i = x_{i_1} \cdots x_{i_s}$ and $(j_1, \ldots, j_s) \in (\mathbb{N} \setminus \{0\})^s$.

Note that our definition of a deformation is different from the ones given at [1, 5]. In particular, our deformations are monomial ideals.

For a given simplicial complex Γ, one can easily find a deformation M of the Stanley-Reisner ideal I_Γ such that M is a generic monomial ideal. In this paper, we characterize simplicial complexes such that their corresponding Stanley-Reisner ideals have deformations that are generic and Cohen-Macaulay. To be precise, we provide algorithms to construct generic Cohen-Macaulay deformations from the Stanley-Reisner ideals of matroids complexes, shifted complexes, and tree complexes. A version of this question appeared in [5] where generic Cohen-Macaulay monomial ideals have been studied extensively.

Definition 1.5. Let $M = \langle m_1, \ldots, m_t\rangle$ be a monomial ideal in S. For $\sigma \subset [t]$, let $m_\sigma := \operatorname{lcm}(m_i : i \in \sigma)$. The *Scarf complex* of M is

$$\Delta_M := \{\sigma \subseteq [t] : m_\sigma \neq m_\tau \text{ for all } \tau \subseteq [t] \text{ and } \tau \neq \sigma\}.$$

For D large enough (larger than any exponent of any variable in any minimal generator of M), let

$$M^* := M + \langle x_1^D, \ldots, x_n^D\rangle.$$

The *extended Scarf complex* of M is the Scarf complex Δ_{M^*} of M^*.

It is easy to see that the induced subcomplex of Δ_{M^*} on the set of generators of M is Δ_M. On the other hand, Miller *et. al* [5] proved that if M is generic, then the induced subcomplex of Δ_{M^*} on the set $\{x_1^D, \ldots, x_n^D\}$ is the simplicial complex $V(M)$, where $I_{V(M)} = \operatorname{rad}(M)$.

Example 1.6. Let Γ is the simplicial complex in Figure 1, and let $M = \langle x_1^2x_2^2x_3^2, x_1x_4, x_2^2x_4^2\rangle$. It is clear that M is a generic deformation of I_Γ (in particular, $\operatorname{rad}(M) = I_\Gamma$ and hence $V(M) = \Gamma$). On the other hand, let $M^* := M + \langle x_1^3, x_2^3, x_3^3, x_4^3\rangle$. Figure 2 gives Δ_M and Δ_{M^*}. The set of facets of Δ_{M^*} is

$$\{\{x_1x_4, x_1^2x_2^2x_3^2, x_1^3, x_2^3\}, \{x_1x_4, x_1^2x_2^2x_3^2, x_1^3, x_3^3\}, \{x_1x_4, x_1^2x_2^2x_3^2, x_2^3, x_3^3\},$$
$$\{x_1x_4, x_2^2x_4^2, x_2^3, x_3^3\}, \{x_1x_4, x_2^2x_4^2, x_3^3, x_4^3\}\}.$$



FIGURE 1. The simplicial complex Γ from Example 1.6

.



FIGURE 2. Δ_M and Δ_{M^*} of M in Example 1.6.

The following theorem is proved as part of [5, Theorem 1.7].

Theorem 1.7. *Let M be a generic monomial ideal. For each facet σ of Δ_{M^*}, let*

$$M_\sigma := \langle x_s^{p_s} \; : \; p_s := \deg_{x_s}(m_\sigma) \textit{ and } p_s < D \rangle.$$

Then

$$M = \bigcap_{\sigma \textit{ is a facet of } \Delta_{M^*}} M_\sigma$$

is a minimal irreducible decomposition of M.

Example 1.6 (*cont.*). By Theorem 1.7, each facet of Δ_{M^*} corresponds to an irreducible component of M, and hence Δ_{M^*} gives a minimal irreducible decomposition of M which implies a minimal primary decomposition of M:

$$\begin{aligned} M &= \langle x_3^2, x_4\rangle \cap \langle x_2^2, x_4\rangle \cap \langle x_1^2, x_4\rangle \cap \langle x_1, x_4^2\rangle \cap \langle x_1, x_2^2\rangle \\ &= \langle x_3^2, x_4\rangle \cap \langle x_2^2, x_4\rangle \cap \langle x_1^2, x_1x_4, x_4^2\rangle \cap \langle x_1, x_2^2\rangle. \end{aligned}$$

Theorem 1.8 ([5, Theorem 2.5])**.** *Let M be a generic monomial ideal. Then M has no embedded associated primes if and only if M is Cohen-Macaulay. In this case, both Δ_M and $V(M)$ are shellable.*

For a given simplicial complex Γ, the theorem above implies that if Γ is not shellable, then there is no generic Cohen-Macaulay monomial ideal M such that $V(M) = \Gamma$. In particular, there is no generic Cohen-Macaulay deformation of I_Γ. The following theorem will play a major role subsequently.

Theorem 1.9. *Let $M \subset S$ be a generic monomial ideal. Then M is Cohen-Macaulay if and only if $\dim(\Delta_M) + \dim(V(M)) = n - 2$.*

Proof. Suppose M is Cohen-Macaulay. By Theorem 1.7, each facet of Δ_{M^*} gives an irreducible component of M. In particular, for any facet σ of Δ_{M^*}, we have

$$|\sigma \cap \{m_1, \ldots, m_t\}| = \operatorname{codim}(M) \quad \text{and} \quad |\sigma \cap \{x_1^D, \ldots, x_n^D\}| = \dim(R/M).$$

Furthermore, both cardinalities are independent of the facet σ, since M has no embedded primes. But $\sigma \cap \{m_1, \ldots, m_t\}$ is a face of Δ_M and any facet of Δ_M is a restriction of a facet of Δ_{M^*}. Thus $\dim(\Delta_M) = \operatorname{codim}(M) - 1$. Hence $\sigma \cap \{x_1^D, \ldots, x_n^D\}$ is a facet of $V(M)$ and therefore $\dim(V(M)) = \dim(R/M) - 1$. But $\operatorname{codim}(M) + \dim(R/M) = n$. Thus, $\dim(\Delta_M) + \dim(V(M)) = n - 2$.

Conversely, suppose $\dim(\Delta_M) + \dim(V(M)) = n - 2$. Then it is enough to show that for any facet σ of Δ_{M^*}, we have $|\sigma \cap \{m_1, \ldots, m_t\}| = n - d$, where $\dim(V(M)) = d - 1$, i.e., $\dim(\Delta_M) = n - d - 1$. This proves that all irreducible components are of the same codimension, namely $n - d$, and hence there are no inclusions among them. Thus the irreducible decomposition is minimal and hence all the associated primes of M are of the same codimension. Thus M has no embedded primes. By Theorem 1.8, M is then Cohen-Macaulay.

To this end, let σ be a facet of Δ_{M^*}. If $|\sigma \cap \{m_1, \ldots, m_t\}| > n - d$, then $\dim(\Delta_M) > n - d - 1$, a contradiction. Now if $|\sigma \cap \{m_1, \ldots, m_t\}| < n - d$, then $|\sigma \cap \{x_1^D, \ldots, x_n^D\}| > d$. Thus $\dim(V(M)) > d - 1$, a contradiction. Therefore, M is Cohen-Macaulay. □

Example 1.10. Let Γ be the 1-dimensional simplicial complex (graph) in Figure 3. It is clear that Γ is shellable, since it is connected. The Stanley-Reisner ideal of Γ is

$$I_\Gamma = \langle x_1x_2x_3, x_1x_4, x_1x_5, x_2x_4, x_3x_5\rangle.$$

Let $M_1 = \langle x_1^3x_2^3x_3^3, x_2^3x_4^3\rangle$, $M_2 = \langle x_1^2x_4^2, x_3^2x_5^2\rangle$, $M_3 = \langle x_1x_5\rangle$, and let $M = M_1 + M_2 + M_3$. It is clear that M is a deformation of I_Γ. Moreover, M is



FIGURE 3. The simplicial complex Γ from Example 1.10.

generic. Namely, for $1 \leq i \leq 2$, given any two distinct minimal generators $m_1, m_2 \in M_i$, there exists a third generator m_3 in M_j for some $j > i$ such that m_3 strictly divides $\operatorname{lcm}(m_1, m_2)$. Thus $\dim(\Delta_M) \leq 2 = 5-2-1 = n-d-1$. But $\{x_1x_5, x_1^2x_4^2, x_2^3x_4^3\}$ is clearly a face of Δ_M. So $\dim(\Delta_M) = 2$. Now, by Theorem 1.9, M is Cohen-Macaulay.

In the next sections, we show that the Stanley-Reisner ideals of matroid, shifted, or tree complexes have generic Cohen-Macaulay deformations.

2. MATROID COMPLEXES

In this section, we prove that the Stanley-Reisner ideals of matroid complexes have generic Cohen-Macaulay deformations. There are many equivalent definitions of matroids, see [6], or [8, §III.3]. The following definition uses the so-called *circuits* axiom.

Definition 2.1. A simplicial complex Δ is a *matroid* if for any two minimal non-faces (circuits) α and β with an $i \in \alpha \cap \beta$, there exists a minimal non-face γ such that $\gamma \subseteq (\alpha \cup \beta) \setminus \{i\}$.

The following corollary follows directly from [8, Proposition 3.1].

Corollary 2.2. *Let* Γ *be a matroid complex on the vertex set* $[n]$*. For every subset* W *of* $[n]$*, the induced subcomplex* $\Gamma_W := \{\sigma \in \Gamma : \sigma \subseteq W\}$ *of* Γ *is a matroid complex. In particular,* Γ_W *is pure and shellable.*

Example 2.3. The simplicial complex Γ in Figure 4 is a matroid. The Stanley-Reisner ideal of Γ is $I_\Gamma = \langle x_1x_3, x_1x_2x_4, x_2x_3x_4 \rangle$.

The following theorem is the main result of this section.

Theorem 2.4. *Let* Γ *be a matroid complex on the vertex set* $[n]$ *and suppose* $\dim(\Gamma) = d-1$*. Then there exists a generic Cohen-Macaulay deformation of* I_Γ*.*

Proof. Pick a facet σ of Γ. Without loss of generality, $\sigma = \{1, \ldots, d\} = [d]$. There exists a unique minimal non-face $\alpha \subset [d+1]$: By Corollary 2.2, the induced subcomplex $\Gamma_{[d+1]}$ is pure and $\sigma \in \Gamma_{[d+1]}$, in particular,



FIGURE 4. The matroid complex Γ from Example 2.3.

$\dim(\Gamma_{[d+1]}) = d-1$. But $\Gamma_{[d+1]}$ is pure, so there exists a unique minimal non-face $\alpha \subset [d+1]$. Let $m_\alpha = \prod_{j\in\alpha} x_j$. Let $M_1 := \langle m_\alpha \rangle$, a monomial ideal in S. By induction, for $i > 1$, let

$$\Sigma_i := \{\alpha \subset [d+i] \, : \, \alpha \text{ is a minimal non-face of } \Gamma \text{ and } d+i \in \alpha\},$$

be the set of minimal non-faces in $[d+i]$ but not in $[d+i-1]$. Let

$$M_i := \langle \prod_{j\in\alpha} x_j^i \, : \, \alpha \in \Sigma_i \rangle.$$

Since we have n vertices, we will have all the minimal non-faces at the inductive step $n-d$.

Let $M = M_1 + \cdots + M_{n-d}$. Then M is generic and $V(M) = \Gamma$. It remains to show that M is Cohen-Macaulay. By Theorem 1.9, it is enough to show that $\dim(\Delta_M) = n-d-1$.

First, we show that, for all $i > 0$, $M_i \neq 0$: at each inductive step i , $2 \leq i \leq n-d$, $\Gamma_{[d+i]}$ is a matroid complex on $d+i$ vertices and $\dim(\Gamma_{[d+i]}) = d-1$. For if $M_i = 0$, there is no missing face with the vertex $d+i$, in particular $\sigma \cup \{d+i\} \in \Gamma$. This is a contradiction, since σ is a facet of Γ. Furthermore, $\dim(\Delta_M) = n-d-1$: For each $1 \leq i \leq n-d$, let

$$F_i := \sigma \cup \{d+i\}.$$

Then there exists a unique minimal non-face (circuit) c_i of Γ such that $d+i \in c_i \subset F_i$. Let

$$m_i := \prod_{j\in c_i} x_j^i.$$

Let $G = \{m_1, \ldots, m_{n-d}\}$. We will show that G is a face of Δ_M and hence $\dim(\Delta_M) = n-d-1$. Notice that

$$\operatorname{lcm}(G) := \operatorname{lcm}(m_1, \ldots, m_{n-d}) = f \cdot x_{d+1} x_{d+2}^2 \cdots x_n^{n-d},$$

where f is a monomial in the variables $x_1, \ldots, x_d$. Moreover, for any proper subset $H \subset G$, it is easy to see that $\operatorname{lcm}(H) \neq \operatorname{lcm}(G)$.

Suppose that, for some $1 \leq t \leq n-d$, there exists a generator $m \in M_t$ such that m divides $\operatorname{lcm}(G)$, say $m = x_{i_1}^t x_{i_2}^t \cdots x_{i_s}^t x_{d+t}^t$, where $1 \leq i_1 \leq$

$\cdots \leq i_s < d+t$. Thus $i_s \leq d$. This implies that m corresponds to the unique minimal non-face $\alpha = \{i_1, \dots, i_s, d+t\}$ which is m_t. Hence G is a face of Δ_M and therefore $\dim(\Delta_M) = n-d-1$. $\square$

Example 2.5. For the simplicial complexes Γ in Figure 4, $\sigma = \{1,2\}$ and $\Gamma_{[3]}$ has the facets σ and $\{2,3\}$. Thus $M_1 = \langle x_1x_3 \rangle$. It is clear that $\Gamma_{[4]} = \Gamma$. Thus we need to compute only M_2. But $\Sigma_2 = \{\{1,2,4\},\{2,3,4\}\}$. Thus $M_2 = \langle x_1^2x_2^2x_4^2, , x_2^2x_3^2x_4^2 \rangle$. Let $M = M_1 + M_2 = \langle x_1x_3, x_1^2x_2^2x_4^2, , x_2^2x_3^2x_4^2 \rangle$. It is clear that M is generic. Moreover, $\dim(\Delta_M) = 1$, see Figure 5. Thus M is Cohen-Macaulay, by Theorem 1.8.



FIGURE 5. The Scarf complex Δ_M of M in Example 2.5.

3. SHIFTED COMPLEXES

In this section, we prove that shellable shifted complexes have generic Cohen-Macaulay deformations.

Definition 3.1. A simplicial complex Γ on the vertex set $[n]$ is *shifted* if, for all $\sigma \in \Gamma$, whenever $j \in \sigma$, $i \in [n]$, and $i < j$, we have $(\sigma \setminus \{j\}) \cup \{i\} \in \Gamma$.

Example 3.2. Let Γ be the 2-dimensional simplicial complex on the vertex set $[6]$ with the set of facets $\{\{1,2,3\},\{1,2,4\},\{1,3,4\},\{1,2,5\},\{1,2,6\}\}$, see Figure 6. It is easy to see that Γ is a shifted complex.



FIGURE 6. The shifted complex Γ from Example 3.2.

Notice that the labeling plays a major role in this definition. So relabeling (permuting the vertices) a shifted complex might yield a non-shifted one, although they are *combinatorially equivalent* (their face lattices are isomorphic).

The following theorem [3, Theorem 3] gives simple geometrical and combinatorial characterizations for Cohen-Macaulay, and shellable shifted complexes.

Theorem 3.3. *Let Γ be a $(d-1)$-dimensional shifted simplicial complex.*

1. Γ *is homotopically equivalent to a wedge of spheres,*
2. Γ *is shellable if and only if* Γ *is Cohen-Macaulay if and only if* Γ *is pure if and only if for every* $\sigma \in \Gamma$ *and* $|\sigma| = t$, $\sigma \cup [d-t] \in \Gamma$.

Let Γ be a shellable $(d-1)$-dimensional shifted simplicial complex on the vertex set $[n]$. We give a partition for the set of minimal non-faces Σ_Γ. Recall that

$$\Sigma_\Gamma := \{\sigma \subseteq [n] \,:\, \sigma \notin \Gamma \text{ but } \sigma \setminus \{i\} \in \Gamma \text{ for each } i \in \sigma\}.$$

Now, for $1 \leq j \leq n-d$, let

$$\Sigma_j := \{\sigma \in \Sigma_\Gamma \,:\, \sigma = A \cup \{d - |A| + j\} \text{ and } A \subseteq \overline{[d-|A|+j]}\}.$$

In the next proposition, we present some facts that will be very useful in proving the main result of this section. But before that we give an example.

Example 3.2 (*cont.*)**.** It is clear that

$$\Sigma_\Gamma = \{\{2,3,4\}, \{3,5\}, \{3,6\}, \{4,5\}, \{4,6\}, \{5,6\}\}.$$

Moreover, it is easy to see that

$$\Sigma_1 = \{\{2,3,4\}, \{3,5\}, \{3,6\}\}.$$
$$\Sigma_2 = \{\{4,5\}, \{4,6\}\}.$$
$$\Sigma_3 = \{\{5,6\}\}.$$

Hence, $\Sigma_\Gamma = \Sigma_1 \cup \Sigma_2 \cup \Sigma_3$.

Proposition 3.4. *Let* Γ, Σ_Γ, *and* Σ_j *be as above.*

1. *For* $1 \leq j \leq n-d$, *we have* $\Sigma_j \neq \emptyset$,
2. *For* $1 \leq i < j \leq n-d$, $\Sigma_i \cap \Sigma_j = \emptyset$, *and*
3. $\Sigma_\Gamma = \cup_{j=1}^{n-d} \Sigma_j$.

Proof. 1. Let $1 \leq j \leq n-d$ and let $H := \{d+j, d+j-1, \ldots, j\}$. Since $|H| = d+1$ and $\dim(\Gamma) = d-1$, for some $j \leq t \leq d+j$, we have that $\{d+j, d+j-1, \ldots, t\}$ is a minimal non-face of Γ. Let $A := \{d+j, d+j-1, \ldots, t-1\}$. It is clear that $|A| = d-t+j$ and hence $t = d-|A|+j$. Therefore, $A \cup \{t\} \in \Sigma_j$ which implies that $\Sigma_j \neq \emptyset$.

2. Straightforward.

3. It is clear that, for any $1 \leq j \leq n-d$, we have $\Sigma_j \subseteq \Sigma_\Gamma$. Let $\sigma \in \Sigma_\Gamma$, say $\sigma = \{i_1, \ldots, i_{t+1}\}$ where $1 \leq i_{t+1} < \cdots < i_1 \leq n$. Let $A := \{i_1, \ldots, i_t\}$.

Since σ is a minimal non-face, we have $A \in \Gamma$. Since $1 \leq i_{t+1} \leq n-t$, there exists $1 \leq j_0 \leq n-d$ such that $i_{t+1} = d-t+j_0$. Notice that $|A| = t$. Thus $A \subseteq \overline{[d-|A|+j_0]}$ and $\sigma = A \cup \{d-|A|+j_0\}$. Therefore $\sigma \in \Sigma_{j_0}$. Hence $\Sigma_\Gamma = \cup_{j=1}^{n-d} \Sigma_j$. □

Proposition 3.5. *Let* Γ, Σ_j *be as above. Let* $\sigma, \gamma \in \Sigma_j$ *and* $\sigma \neq \gamma$. *There exists* $s > j$ *and* $\tau \in \Sigma_s$ *such that* $\tau \subset \sigma \cup \gamma$.

Proof. Since $\sigma, \gamma \in \Sigma_j$, there exists $A, B \in \Gamma$ such that $\sigma = A \cup \{d-|A|+j\}$ and $\gamma = B \cup \{d-|B|+j\}$ where $A \subseteq \overline{[d-|A|+j]}$ and $B \subseteq \overline{[d-|B|+j]}$. Moreover, $A \neq B$, since $\sigma \neq \gamma$. Without loss of generality, assume $|A| \leq |B|$.
Case 1. Suppose $b \leq d-|A|+j$, for every $b \in B \setminus A$. Thus

$$B \setminus A \subseteq \{d-|A|+j, \ldots, d-|B|+j+1\}.$$

Hence

$$\begin{aligned} |B \setminus A| &\leq \{d-|A|+j, \ldots, d-|B|+j+1\}| \\ &= (d-|A|+j) - (d-|B|+j) \\ &= |B| - |A|, \end{aligned}$$

which implies that $A \subset B$. But $d-|A|+j \notin B$, otherwise $\sigma \subset B$. Therefore,

$$B \subseteq A \cup \{d-|A|+j-1, \ldots, d-|B|+j+1\}.$$

Hence

$$\begin{aligned} |B| &\leq |A \cup \{d-|A|+j, \ldots, d-|B|+j+1\}| \\ &= |A| + (d-|A|+j-1) - (d-|B|+j) \\ &= |B| - 1, \end{aligned}$$

a contradiction.
Case 2. There exists $b \in B \setminus A$ such that $b > d-|A|+j$, say $b = d-|A|+j+h$ for some $h \geq 1$. Since σ is a non-face and $b > d-|A|+j$, the set $A \cup \{b\}$ must contain a minimal non-face. Let

$$\begin{aligned} A^> &:= \{a \in A \,:\, a > b\} = \{a_1 > \cdots > a_t\}, \text{ and} \\ A^< &:= \{a \in A \,:\, a < b\} = \{b_1 > \cdots > b_{|A|-t}\}, \end{aligned}$$

if $A^> = \emptyset$, let $t := 0$. If $A^< \neq \emptyset$, then for $1 \leq i \leq |A|-t$,

$$b_i = d-|A|+j+s_i,$$

where $s_i \geq |A|-t-i+1$. Thus

$$h > s_1 > \cdots > s_{|A|-t} \geq 1.$$

In the following, we will discuss all possible cases.
1. Suppose $A^> \cup \{b\}$ is a minimal non-face of Γ. We need to show that $A^> \cup \{b\} \in \Sigma_s$ for some $s > j$. So it is enough to show that $b = d-|A^>|+s =$

$d-t+s$, for some $s>j$. But

$$\begin{aligned} b > b_1 &= d-|A|+j+s_1 \\ &\geq d-|A|+j+|A|-t \\ &= d-|A^>|+j. \end{aligned}$$

This implies that $b = d-|A^>|+s$ for some $s>j$. Therefore, $A^> \cup \{b\} \in \Sigma_s$ and $s>j$.

2. Suppose $A^> \cup \{b\} \cup \{b_1,\dots,b_i\}$ is a minimal non-face of Γ. As above,

$$\begin{aligned} b_i &= d-|A|+j+s_i \\ &\geq d-|A|+j+|A|-t-i+1 \\ &= d-(t+i)+j+1. \end{aligned}$$

Thus $b_i = d - |A^> \cup \{b\} \cup \{b_1,\dots,b_{i-1}\}| + s$ for some $s>j$. Therefore,

$$(A^> \cup \{b\} \cup \{b_1,\dots,b_{i-1}\}) \cup \{b_i\} \in \Sigma_s.$$

$\square$

In the following we construct a generic deformation of I_Γ. For $1 \leq j \leq n-d$, let

$$M_j := \langle \prod_{i\in\sigma} x_i^{n-d-j+1} \; : \; \sigma \in \Sigma_j \rangle,$$

a monomial ideal in $R := k[x_1,\dots,x_n]$. Let

$$M := M_1 + \dots + M_{n-d}.$$

The following corollary is straightforward from Proposition 3.5.

Corollary 3.6. *For any $1 \leq j < n-d$, let $m_1, m_2 \in M_j$ be two distinct generators of M_j. Then there exists $s > j$ and a minimal generator $m_3 \in M_s$ such that m_3 divides* $\operatorname{lcm}(m_1,m_2)$. *Thus M is a generic deformation of I_Γ.*

The following theorem shows that M is Cohen-Macaulay. Before we prove that, we verify it for our Example.

Example 3.2 (*cont.*). The above construction yield the following monomial ideals:

$$\begin{aligned} M_1 &= \langle x_2^3x_3^3x_4^3, x_3^3x_5^3, x_3^3x_6^3 \rangle, \\ M_2 &= \langle x_4^2x_5^2, x_4^2x_6^2 \rangle, \\ M_3 &= \langle x_5x_6 \rangle. \end{aligned}$$

It is easy to check that the monomial ideal

$$M = \langle x_2^3x_3^3x_4^3, x_3^3x_5^3, x_3^3x_6^3, x_4^2x_5^2, x_4^2x_6^2, x_5x_6 \rangle$$

is generic and $\operatorname{rad}(M) = I_\Gamma$. Moreover, by Corollary 3.6, $\dim(\Delta_M) \leq 2$. To show that M is Cohen-Macaulay, it is enough to show that $\dim(\Delta_M) = 2$, by Theorem 1.9. But it is clear that $\operatorname{lcm}\{x_5x_6, x_4^2x_5^2, x_3^3x_5^3\}$ is uniquely attained and hence $\{x_5x_6, x_4^2x_5^2, x_3^3x_5^3\}$ corresponds to a facet of Δ_M. Therefore $\dim(\Delta_M) = 2$ and hence M is Cohen-Macaulay.

Theorem 3.7. *The ideal M, constructed above, is a generic Cohen-Macaulay integral deformation of I_Γ.*

Proof. By Corollary 3.6, M is a generic integral deformation of I_Γ. So it remains to show that M is Cohen-Macaulay. By Theorem 1.9, it is enough to show that $\dim(\Delta_M) = n-d-1$. By Corollary 3.6, $\dim(\Delta_M) \leq n-d-1$. Thus to show that $\dim(\Delta_M) = n-d-1$, it is enough to find a face F of Δ_M such that $\dim(F) = n-d-1$.

For $1 \leq j \leq n-d$, let

$$\sigma_j := \{d+j, d+j-1, \dots, i_j\},$$

where i_j is small enough such that $\sigma_j \in \Sigma_\Gamma$, i.e., σ_j is a minimal non-face of Γ. Such an i_j exists since $|[d+j]| > d$ for $j > 1$ and $\dim(\Gamma) = d-1$. First we will show that $\sigma_j \in \Sigma_j$. Let $A := \sigma_j \setminus \{i_j\}$. It is clear that $|A| = (d+j) - i_j$. Thus $i_j = d - |A| + j$. Therefore $A \subsetneq \overline{[d-|A|+j]}$ and hence $\sigma = A \cup \{i_j\} \in \sigma_j$.

Next we will show that $F := \{m_{\sigma_1}, \dots, m_{\sigma_{n-d}}\}$ is a face of Δ_M. Now

$$\begin{aligned} \operatorname{lcm}(F) &= \operatorname{lcm}(m_{\sigma_1}, \dots, m_{\sigma_{n-d}}) \\ &= x_n x_{n-1}^2 \cdots x_{d+1}^{n-d} x_d^{n-d} \cdots x_{i_1}^{n-d}. \end{aligned}$$

It is clear that, for $1 \leq j \leq n-d$, we have $\operatorname{lcm}(F) \neq \operatorname{lcm}(F \setminus \{m_{\sigma_j}\})$. So we only need to show that $\operatorname{lcm}(F) \neq \operatorname{lcm}(F \cup \{m_\sigma\})$ for any $\sigma \in \Sigma_\Gamma \setminus F$. Let $\sigma \in \Sigma_\Gamma \setminus F$, say $\sigma \in \Sigma_j$ for some $1 \leq j \leq n-d$. Thus $\sigma = A \cup \{d-|A|+j\}$, for some $A \in \Gamma$ and $A \subseteq \overline{[d-|A|+j]}$. Since $\sigma \notin F$, there exists $a \in A$ such that $a > d+j$, otherwise $\sigma = \sigma_j$. Suppose $a = d+i$ for some $i > j$. Thus $a \in \sigma_i$ and hence $\deg_{x_a}(\operatorname{lcm}(F)) = n-(d+i)+1$. But $\deg_{x_a}(m_\sigma) = n-(d+j)+1$. Therefore, $\operatorname{lcm}(F) \leq \operatorname{lcm}(F \cup \{m_\sigma\})$. Hence F is a face of Δ_M of dimension $n-d-1$. Therefore, $\dim(\Delta_M) = n-d-1$. By Theorem 1.9, M is Cohen-Macaulay. □

4. Tree Complexes: Shellable Clique Complexes of Chordal Graphs

In this section, we study the class of shellable clique complexes of chordal graphs, which we call tree complexes because they generalize tree graphs. The main result of this section is that tree complexes have generic Cohen-Macaulay deformations. Moreover, we give an exact formula to compute the f-vector of tree complexes.

Throughout this section, let G be a simple graph (undirected graph with no loops and no multiple edges) on the vertex set $[n]$.

Definition 4.1. Let G be as above.

(1) A subset $\sigma \subseteq [n]$ is a *clique* if the induced subgraph G_σ is complete.

(2) The *clique complex* of G is

$$K(G) := \{\sigma \subseteq [n] : \sigma \text{ is a clique of } G\}.$$

Example 4.2. A graph G and its clique complex $K(G)$ are in Figure 7. Notice that G is the 1-skeleton of $K(G)$.



FIGURE 7. A graph G and its clique complex $K(G)$.

The proof of the following lemma is straightforward.

Lemma 4.3. *Every minimal non-face of $K(G)$ has precisely two elements. Moreover, if Δ is a simplicial complex such that every minimal non-face has exactly two elements, then $\Delta = K(G)$, where G is the 1-skeleton of Δ.*

Clique complexes are also called *flag* complexes, see [9].

Definition 4.4. A simplicial complex Δ is a *tree* complex if there exists a shelling $F_1, \ldots, F_t$ of Δ such that, for $1 < i \le t$, there exists $j_i \in F_i$ such that $j_i \notin F_1 \cup \cdots \cup F_{i-1}$. In this case, we say that F_i introduces the vertex j_i.

It is easy to see that any tree graph is a 1-dimensional tree complex. Hence tree complexes are a natural generalization of tree graphs.

Example 4.5. The simplicial complex Γ in Figure 8 is a 2-dimensional tree complex.

One of the most interesting problems in combinatorics is characterizing the f-vectors of flag complexes [2, 9, 4]. For tree complexes, the next theorem gives an exact formula for the f-vector of these complexes.

Theorem 4.6. *Let Δ be a $(d-1)$-dimensional tree complex on $[n]$. Then the number of facets of Δ is $f_{d-1}(\Delta) = n - d + 1$, and for $1 < i \le d - 1$,*

$$\begin{aligned} f_{i-1}(\Delta) &:= \binom{d}{i} + (n-d)\{\binom{d}{i} - \binom{d-1}{i}\} \\ &= (n-d+1)\binom{d}{i} - (n-d)\binom{d-1}{i}. \end{aligned}$$

Proof. Let $F_1, \ldots, F_t$ be a shelling of Δ. By Theorem 4.10, every vertex $d \le i \le n$ introduces a facet. Thus $f_{d-1}(\Delta) = n - d + 1$. Now for $1 < i \le d - 1$, it is clear that F_1 has $\binom{d}{i}$ $(i-1)$-faces. Moreover, every facet



Figure 8. The tree complex from Example 4.5.

F_j, where $j > 1$ has $\binom{d}{i}$ $(i-1)$-faces minus the the number of $(i-1)$-faces that F_j shares with $F_1, \ldots, F_{j-1}$. But F_j shares only $d-1$ vertices with $F_1, \ldots, F_{j-1}$. Hence the number of $(i-1)$-faces it shares is $\binom{d-1}{i}$ which implies the equality above. □

Definition 4.7. A graph G is *chordal* if every cycle of length four or more has a chord (an edge linking two non-adjacent nodes in the cycle).

Example 4.8. The graph G in Figure 7 is chordal.

The following theorem gives many useful characterizations of chordal graphs.

Theorem 4.9 ([7, Theorem 7, p 112]). *The following statements are equivalent.*

(1) *G is a chordal graph.*

(2) *All vertices of G can be deleted by arranging them in separate piles, one for each maximal clique, and then repeatedly applying the following two operations:*
  - *Delete a vertex that occurs in only one pile.*
  - *Delete a pile if all its vertices appear in another pile.*

(3) *There is a spanning tree (called a clique tree) T of the facet graph of $K(G)$ such that for every vertex i of G, if we remove from T all cliques not containing i, the remaining subtree stays connected. In other words, any two cliques containing i are either adjacent in T or connected by a path made entirely of cliques that contain i.*

Suppose Δ is a $(d-1)$-dimensional tree complex on $[n]$. Without loss of generality (after relabeling the vertices), we may assume $F_1 := [d]$ and, for $1 < i \leq t$, F_i introduces the vertex $d+i-1$, i.e., $F_i := A \cup \{d+i-1\}$ for some $A \subseteq [d+i-2]$ and $|A| = d-1$. The following theorem shows that the class of tree complexes is equal to the class of shellable clique complexes of chordal graphs.

Theorem 4.10. *A shellable simplicial complex Δ is a tree complex if and only if $\Delta = K(G)$, the clique complex of a chordal graph G.*

Proof. Let $F_1, \ldots, F_t$ be a shelling of Δ. First suppose that $\Delta = K(G)$, for some chordal graph G, and assume that Δ is not a tree complex. Let s be the smallest index such that F_s does not introduce a new vertex. There exists a simple cycle in the facet graph of $K(G)$ that contains F_s, without loss of generality, $F_1, \ldots, F_s$ is a simple cycle of minimal length. Thus every vertex is contained in more than one clique. This is a contradiction to (2) of Theorem 4.9. Therefore, for $1 < i \leq t$, there exists $i_j \in F_i$ such that $i_j \notin F_1 \cup \cdots \cup F_{i_1}$ and hence Δ is a tree complex.

Now suppose that Δ is a tree complex. Let G be the 1-skeleton of Δ (G is the subcomplex of simplexes of dimension less than or equal to 1). It is clear that G is a graph on the set of vertices $[n]$. Indeed G is a chordal graph: For every facet F_i, let P_i be the pile that contains the vertices of F_i. Since, for $1 < i \leq t$, F_i introduces the vertex j_i, it is clear that one can empty all the piles by deleting each vertex that is in only one pile and then deleting any pile where its vertices are in another pile. Thus G is chordal. So we only need to show that $\Delta = K(G)$. It is clear that Δ is a subcomplex of $K(G)$. Suppose C is a clique of G and $h \in C$ is the maximal element. Thus the vertex h is introduced by the facet F_{i_h}, for some $1 \leq i_h \leq t$. Thus every vertex connected to h and less than h is in F_{i_h} which implies that $C \subset F_{i_h}$. Hence $\Delta = K(G)$. ☐

Let $FG(\Delta)$ be the facet graph of Δ. By Theorem 4.6, $FG(\Delta)$ has $n-d+1$ facets. Let T_Δ be a clique tree of $FG(\Delta)$, it has $n-d$ edges and each edge in T_Δ corresponds to a missing edge in Δ:

If F_i and F_j are vertices connected by an edge e in T_Δ, then F_i and F_j are facets of Δ, i.e., $|F_i| = |F_j| = d$ and $|F_i \cap F_j| = d-1$. Thus, since Δ is a clique complex, there is a unique minimal non-face in $F_i \cup F_j$, say $\{p_e, q_e\}$. Let

$$\Sigma_0 := \{\{p_e, q_e\} \,:\, e \in T_\Delta\}.$$

Now for the edge e, let

$$F_{ij} := F_i \cup F_j.$$

Let Δ_1 be the d-dimensional simplicial complex on $[n]$ with the set of facets $\{F_{ij} \,:\, e \in T_\Delta\}$. It is easy to see that Δ_1 is a tree complex. If Δ_1 is not a simplex, let T_{Δ_1} be a clique tree of $FG(\Delta_1)$. Repeat the steps above for this case (Δ_t is a simplex only when $t = n - d$). Hence, in general, for $1 \leq t < n - d$, let

$$\Sigma_t := \{\{p_e, q_e\} \,:\, e \in T_{\Delta_t}\}.$$

This process gives a partition of Σ_Γ. Before we give a proof of this fact, we will verify it for our example.

Example 4.5 (*cont.*). It is easy to check that

$$\Sigma_\Gamma = \{\{3,5\}, \{4,5\}, \{1,6\}, \{2,7\}, \{3,4\}, \{4,6\}, \{1,7\}, \{3,6\}, \{4,7\}, \{3,7\}\}.$$

Moreover, the graph in Figure 8 (i) is $FG(\Gamma_i)$ and the subgraph of dashed edges is T_{Γ_i}. This yields the following partition of Σ_Γ.

$$\begin{aligned}
\Sigma_0 &= \{\{3,5\}, \{4,5\}, \{1,6\}, \{2,7\}\}\\
\Sigma_1 &= \{\{3,4\}, \{4,6\}, \{1,7\}, \{3,6\}, \{4,7\}\}\\
\Sigma_2 &= \{\{3,6\}, \{4,7\}\}\\
\Sigma_3 &= \{\{3,7\}\}.
\end{aligned}$$

The proof of the following proposition is straightforward.

Proposition 4.11. *Let Δ be a tree complex and Σ_t be as above.*

(1) *For $0 \leq t < n - d$, we have $\Sigma_t \neq \emptyset$,*
(2) *For $0 \leq t_1 < t_2 < n - d$, $\Sigma_{t_1} \cap \Sigma_{t_2} = \emptyset$, and*
(3) *$\Sigma_\Delta = \cup_{t=0}^{n-d-1} \Sigma_t$.*

The following proposition will help us to construct a generic deformation from I_Δ.

Proposition 4.12. *Let Δ, Σ_t be as above. Let $e_1, e_2 \in \Sigma_t$ and $e_1 \neq e_2$. There exists $s > t$ and $e_3 \in \Sigma_s$ such that $e_3 \subset e_1 \cup e_2$.*

Proof. Without loss of generality, let

$$e_1 = \{p_1, q_1\}, \qquad e_2 = \{p_2, q_2\}$$

be two edges of the clique tree T_{Δ_t} where the edge e_1 (resp. e_2) has as vertices the $(d + t - 1)$-simplexes (facets) τ_1, τ_2 (resp. γ_1, γ_2) of Δ_t, say

$$\begin{aligned}
\tau_1 &= A \cup \{p_1\}, & \tau_2 &= A \cup \{q_1\},\\
\gamma_1 &= B \cup \{p_2\}, & \gamma_2 &= B \cup \{q_2\}.
\end{aligned}$$

Case 1. $p := p_1 = p_2$ ($q_1 \neq q_2$). If $\tau_1 = \gamma_1$, then $A = B$. Thus the edge $\{q_1, q_2\}$ is in the facet graph $FG(\Delta_t)$ but not in T_{Δ_t}, otherwise

τ_1, τ_2, γ_2 form a cycle in T_{Δ_t}. So assume that $\tau_1 \neq \gamma_1$. There exists a path $\tau_1 = G_1, \ldots, G_r = \gamma_1$ such that $p \in G_i$ for $1 \leq i \leq r$. This implies that $\tau_2, G_1, \ldots, G_r, \gamma_2$ is the only path between τ_2 and γ_2 in T_{Δ_t}. Hence $\{q_1, q_2\}$ is not an edge in T_{Δ_t}, otherwise we have a cycle in T_{Δ_t}. In this case, let $e_3 = \{q_1, q_2\}$.

Case 2. $p_i \neq q_j$ for $1 \leq i, j \leq 2$. Suppose for $1 \leq i, j \leq 2$, the edge $\{p_i, q_j\} \in \Delta_t$. Thus p_1, q_1, p_2, q_2 form a 4-cycle in the chordal graph G, hence either the edge $\{p_1, q_1\}$ or $\{p_2, q_2\}$ is in G, a contradiction. Without loss of generality, suppose the 1-simplex $\{p_1, p_2\} \notin \Delta_t$. If $\{p_1, p_2\} \notin T_{\Delta_t}$, then we are done. Suppose $\{p_1, p_2\}$ is an edge in T_{Δ_t} with the vertices F and G such that $p_1 \in F$ and $p_2 \in G$. Since $p_1 \in F \cap \tau_1$, there exists a path between τ_1 and F made entirely from facets containing p_1. Similarly, there exists a path between G and γ_1 made entirely from facets containing p_2. Thus there is a path between τ_2 and γ_2. It is clear that if $\{q_1, q_2\} \in \Delta_t$ or $\{p_1, p_2\}$ is an edge in T_{Δ_t}, then we have a cycle in T_{Δ_t}. Once again, in this case, let $e_3 = \{q_1, q_2\}$. □

In the following we construct a generic integral deformation of I_Δ. For $0 \leq t < n - d$, and $\{p_e, q_e\} \in \Sigma_t$, let

$$M_t := \langle x_{p_e}^{n-d-t} x_{q_e}^{n-d-t} : \{p_e, q_e\} \in \Sigma_t \rangle.$$

It is clear that $M_t \neq 0$, otherwise there are no missing faces and hence we have a simplex which is not the case unless $t = n - d$.

Let $M = M_0 + M_1 + \cdots + M_{d-t-1}$. It is clear that $\mathrm{rad}(M) = I_\Delta$. Moreover, in the next Lemma, we show that M is generic. Before that we continue with our example.

Example 4.5 (*cont.*). For each i; $0 \leq i \leq 4$, the corresponding ideals to the partition above are:

$$\begin{aligned}
M_0 &= \langle x_3^4 x_5^4, x_4^4 x_5^4, x_1^4 x_6^4, x_2^4 x_7^4 \rangle, \\
M_1 &= \langle x_3^3 x_4^3, x_4^3 x_6^4, x_1^3 x_7^3 \rangle, \\
M_2 &= \langle x_3^2 x_6^2, x_4^2 x_7^2 \rangle, \\
M_3 &= \langle x_3 x_7 \rangle.
\end{aligned}$$

Since Γ_4 is a simplex, the facet graph $FG(\Gamma_4)$ is a vertex with no edges and hence $M_4 = 0$.

Let $M = M_0 + \cdots + M_3$. It is easy to see that M is a generic monomial ideal and $V(M) = \Gamma$.

The following lemma follows from Corollary 4.12.

Lemma 4.13. *For any $0 \leq j < n - d$, let $m_1, m_2 \in M_j$ be two distinct generators of M_j. Then there exists $s > j$ and a minimal generator $m_3 \in M_s$ such that m_3 divides $\mathrm{lcm}(m_1, m_2)$. Thus M is a generic integral deformation of I_Δ.*

The following theorem shows that M is Cohen-Macaulay. Before we prove that, we verify it for our example.

Example 4.5 (*cont.*). It is easy to see that $\{x_3x_7, x_3^2x_6^2, x_3^3x_4^3, x_3^4x_5^4\}$ is a face of Δ_M and hence $\dim(\Delta_M) = 3 = 7 - 3 - 1$. Thus, by Theorem 1.9, M is Cohen-Macaulay.

Theorem 4.14. *The ideal M is a generic Cohen-Macaulay integral deformation of I_Δ.*

Proof. It remains to show that M is Cohen-Macaulay. Since $V(M) = \Delta$, by Theorem 1.9, we need to show that $\dim(\Delta_M) = n - d - 1$. By Lemma 4.13, each face of Δ_M is of dimension $\leq n - d - 1$. To show that $\dim(\Delta_M) = n - d - 1$, it is enough to find a face of dimension $n - d - 1$. For $0 \leq i < n - d$, there exists a unique minimal non-face in Δ_i which is a subset of $[d + i + 1]$ and contains $d + i + 1$, say $\{\alpha_i, d + i + 1\} \in \Sigma_i$. Let $m_i = x_{\alpha_i}^{n-d-i} x_{d+i+1}^{n-d-i}$ be the corresponding monomial in M_i.
Claim. $C := \{m_0, \ldots, m_{n-d-1}\}$ is a facet of Δ_M and hence $\dim(\Delta_M) = n - d - 1$.

$$\mathrm{lcm}(C) := \mathrm{lcm}\{m_0, \ldots, m_{n-d-1}\} = f \cdot x_{d+1}^{n-d} x_{d+2}^{n-d-1} \cdots x_{d+i}^{n-d-i+1} \cdots x_{n-1}^2 x_n,$$

where f is a monomial on the variables $x_1, \ldots, x_d$. It is clear that $\mathrm{lcm}(C) \neq \mathrm{lcm}(H)$ for any proper subset $H \subset C$. Suppose there exists i, j such that $j < d + i + 1$, $\{j, d + i + 1\} \in \Sigma_\Delta$ and $x_j^a x_{d+i+1}^a$ divides $\mathrm{lcm}(C)$, i.e., $a \leq n - d - i + 2$. This implies that $\{j, d + i + 1\}$ corresponds to an edge in Δ_s where $s \geq i$. Since $\{\alpha_e, d + i + 1\}$ is the only non-face in Δ_i that is a subset of $[d + i + 1]$, we must have $s = i$ and $j = \alpha_e$ and hence $m = m_i$. Therefore, $C \in \Delta_M$ and hence $\dim(\Delta_M) = n - d - 1$. By Theorem 1.9, M is Cohen-Macaulay. ☐

References

[1] D. Bayer, I. Peeva, and B. Sturmfels. Monomial ideals. *Math. Research Letters*, 5:31–46, 1998.

[2] L. Billera and A. Björner. Face numbers of polytopes and complexes. In J. Goodman and J. O'Rourke, editors, *Discrete and Computational Geometry*, pages 291–310. CRC Press, New York, 1997.

[3] A. Björner and G. Kalai. On f-vectors and homology. *Annals New York Academy of Sciences*, 555:63–80, 1989.

[4] G. Kalai. Combinatorics with a geometric flavor: Some examples. preprint.

[5] E. Miller, B. Sturmfels, and K. Yanagawa. Generic and cogeneric monomial ideals. *J. Symbolic Comp.*, 29:691–708, 2000.

[6] J. Oxley. *Matroid Theory.* Oxford University Press, Oxford, 1992.

[7] J. Pearl. *Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference.* Morgan Kaufmann Publications, San Mateo, CA, 1988.

[8] R. Stanley. *Combinatorics and Commutative Algebra*, volume 41 of *Progress in Mathematics.* Birkhäuser, Boston, MA, 2 edition, 1996.

[9] R. Stanley. Positivity problems and conjectures in algebraic combinatorics. In V. Arnold, M. Atiyah, P. Lax, and B. Mazur, editors, *Mathematics: Frontiers and Perspectives*, pages 295–319. American Mathematical Society, Providence, RI, 2000.

[10] K. Yanagawa. f_Δ type free resolutions of monomial ideals. *Proc. Amer. Math. Soc.*, 127:377–383, 1998.

(Abdul Salam Jarrah) Department of Mathematics, East Tennessee State University, Johnson City, TN 37614
E-mail address: `jarrah@mail.etsu.edu`

(Reinhard Laubenbacher) Virginia Bioinformatics Institute, 1880 Pratt Drive, Blacksburg, VA 24061
E-mail address: `reinhard@vbi.vt.edu`